

Main Physical Aspects of the Mathematical Conception of Energy in Thermodynamics

V. P. Maslov

National Research University Higher School of Economics, Moscow, 123458, Russia; Department of Physics, Lomonosov Moscow State University, Moscow, 119234, Russia; A.Ishlinsky Institute for Problems in Mechanics of the Russian Academy of Sciences, 101, prospect Vernadskogo, Moscow, 119526, Russia; v.p.maslov@mail.ru

Abstract

We consider the main physical notions and phenomena described by the author in his mathematical theory of thermodynamics. The new mathematical model yields the equation of state for a wide class of classical gases consisting of non-polar molecules provided that the spinodal, the critical isochore and the second virial coefficient are given. As an example, the spinodal, the critical isochore and the second virial coefficient are taken from the Van-der-Waals model. For this specific example, the isotherms constructed on the basis of the author's model are compared to the Van-der-Waals isotherms, obtained from completely different considerations.

Keywords: Van-der-Waals model; compressibility factor; spinodal; number of degrees of freedom; admissible size of clusters; quasi-static process.

1 Introduction

In the present paper, we present the foundations of the new mathematical model of thermodynamics for the values of the energy for which the molecules do not split into ions and are in the pre-plasma state.

The new mathematical model constructed by the author in the cycle of papers [1]–[2], differs somewhat from the commonly accepted model of phenomenological thermodynamics (see Section 10) and allows us to construct the equation of state for a wider class of classical gases.

We mainly study the metastable states in the case of gases and in the case of fluids up to the critical isochore $\rho = \rho_c$. The liquid isotherms are divided into two parts: the region with temperatures near the critical temperature T_c and the region of the “hard” liquid whose isotherms pass through the point of zero pressure.

Further, the distributions in the Van-der-Waals model are compared to the distributions appearing in the author's new model for classical gases with given critical temperatures, in particular, with Van-der-Waals critical temperature. The precision with which the isotherms constructed from the author's theory coincide with the Van-der-Waals isotherms turns out to be less than the accepted precision in experimental studies. The probability of such a coincidence being accidental is “infinitely small”.

All the distributions of energy levels for classical gas considered in the present paper have been established mathematically, see [1]–[3] and references therein) and have met

with approval (see, for example, the review [4], published in the journal “Probability Theory and its Applications” in connection with the State Prize of the Russian Federation in Science and Technology granted to the author in 2013 (for developing the mathematical foundations of modern thermodynamics).

2 Van-der-Waals normalization and the law of corresponding states

The correct choice of units of measurement for different gases allowed Van-der-Waals to compare the parameters of the gases to dimensionless quantities, thus obtaining the famous “law of corresponding states”.

For every gas, there exists a critical temperature T_c and a critical pressure P_c such that if the temperature or the pressure is greater than their critical values, then gas and liquid can no longer be distinguished. This state of matter is known as a fluid.

The Van-der-Waals “normalization” consists in taking the ratio of the parameters by their critical value, so that we consider the reduced temperature $T_r = T/T_c$ and the reduced pressure $P_r = P/P_c$. In these coordinates, the diagrams of the state of various gases resemble each other – this is a manifestation of the law of corresponding states.

The equation of state for the Van-der-Waals gas in dimensionless variables has the form:

$$(P_r + \alpha\rho^2)(1 - \tilde{\alpha}\rho) = \rho T_r, \quad (1)$$

where $\alpha = 3^3/2^6$, $\tilde{\alpha} = 1/2^3$, and the dimensionless concentration (density) $\rho = N/V$ is chosen so that $\rho_c = 8/3$ in the critical point.

In thermodynamics, the dimension of the internal energy E is equal to the dimension of the product PV of the pressure by the volume. On the other hand, temperature plays the role of mean energy E/N of the gas (or of the liquid). For that reason, when the dimension of temperature is given in energy units, the parameter $Z = P/\rho T$, known as the *compressibility factor*, is dimensionless. For the equation of state of the gas or liquid, the most natural diagram is the P_r - Z diagram, in which the dimensionless pressure P_r is plotted along the x -axis and the dimensionless parameter Z , along the y -axis. A diagram with axes P_r, Z is called a *Hougen–Watson diagram*. In the Van-der-Waals model, the critical value of the compressibility factor is

$$Z_c = \frac{P_c V_c}{N T_c} = \frac{3}{8}.$$

On Fig. 2–1, the Van-der-Waals isotherms are shown in a P_r - Z Hougen–Watson diagram, for which we used the normalization $P_r = P/P_c$, $T_r = T/T_c$, where T_c is the critical temperature.

3 On the number of degrees of freedom and the “effect” of density

In the physics literature, the notion of number of degrees of freedom is used for ideal gases, in which particles do not interact. In probability, the number of degrees of freedom is also considered independently of any interaction.




Figure 1: Van-der-Waals isotherms on the P_r - Z diagram. On the plot $P = P_r$, $T = T_r$. The curve shown by the fat hashed line is the critical isotherm $T_r = 1$. The dotted line is the spinodal, the thin hashed line is the binodal.




Figure 2: The P_r - Z diagram for the Van-der-Waals equation with $P = P_r$.

When physicists speak of the number of degrees of freedom, they usually have in mind the number of degrees of freedom of a single molecule. Normally, a one-atom molecule will have 3 degrees of freedom, a two-atom molecule, 5. In a gas, the molecules move with different speeds and different energies; although their mean energy – temperature – is the same, their individual energies may be different, and the number of degrees of freedom of different molecules can also differ. Thus a two-atom molecule with very high energy may have a number of degrees of freedom greater than 5, and this will affect the mean value (average over all molecules) of the number of degrees of freedom of the given gas. This mean value is called the *collective number of degrees of freedom* of the gas, and is a fractional rather than a whole number.

A one-atom molecule has 3 degrees of freedom, a two-atom molecule, 5. Two-atom molecules are regarded in [5] as molecules of ideal gas. Nevertheless, saying that the number of degrees of freedom of a molecule is 5 means that we are implementing an exclusion rule: we are saying that one of the degrees of freedom is excluded. More precisely, we regard the molecule as a dumbbell, and exclude its oscillations as a rod.

However, if the temperature increases, these “rod” oscillations will take place and so, for sufficiently high temperatures (energies), the molecules will have a greater fractional number of degrees of freedom.

In the author’s model, the number of degrees of freedom is an important independent parameter, so that the notion of exclusion need not be defined by means of some internal considerations ¹.

Although the collective number of degrees of freedom of a classical gas can be determined by the initial interaction between its particles, in the probabilistic description of the generalized ideal gas (a notion that we present in Section 5), this parameter is introduced independently.

If nuclear forces are taken into consideration, it can be shown that a wide class of two-atom molecules have 5 degrees of freedom, but this is difficult to establish. And so we prefer to stipulate axiomatically, as in probability theory, that on a given energy level there can be only one or, say, no more than K particles.

Another important consideration is that we do not take into account the number of particles and the volume separately, we consider their ratio – density (concentration).

The consideration of density alone leads to the following important consequence in thermodynamics: density does not depend on the numeration of particles contained in the given volume. Whatever the numeration of the particles, the density remains the same. It is commonly believed (in particular, it is stated in the book [7]) that classical particles differ from quantum particles in that they can be numbered and then the motion of any individual particle can be followed by keeping track of its number. This is correct, but if the behavior of a multi-particle system is described by equations involving density, e.g. via a probability distribution, then such a description does not depend on the numbering of particles.

Therefore, all the results from quantum mechanics that follow from the remarkable fact that the solution does not depend on numeration can be carried over to the description of any classical multi-particle system by means of equations containing density².

The exclusions that we impose will lead us to a more general distribution in which we stipulate that no more than K particles are on the same energy level. The natural number K will be called the *maximal occupation number* or *maximal admissible size of clusters*.

The corresponding energy distribution, often called parastatistical [9]–[10], will be described in the next section.

Remark 1. In the Landau–Lifshits textbook [5], two relations are given as the basic equations

$$\begin{aligned}\sum_k N_k &= N, \\ \sum_k \varepsilon_k N_k &= \tilde{E},\end{aligned}\tag{2}$$

¹Sometimes, instead of the number of degrees of freedom, such notions as free will or freedom of choice and so on are considered (see, e.g. [6]).

²In particular, the most important thing in the chapter on identical particles in [7] is the method of secondary quantization introduced by Dirac and neatly explained in mathematical terms by Fock (the Fock space, the creation and annihilation operators and so on). Since density does not depend on numeration, the method of secondary quantization had been carried over to classical multi-particle systems described by equations depending on density. This was done by Schoenberg in 1953–54. Later Schoenberg’s approach was generalized in the author’s joint article with O.Yu. Shvedov [8].

where $\tilde{E}$ is the energy, N is the number of particles and $\{\varepsilon_k\}$ is a discrete family of energy levels (see [11], [12]).

The values ε_k in equation (2) are related to the existing interaction potential for the particles. It turns out that if we use the interaction with collective number of degrees of freedom equal to D , then we must put

$$\varepsilon_k = \text{const} \cdot k^{D/2}, \quad (3)$$

i.e.,

$$\tilde{E} = \sum_k \text{const} \cdot k^{D/2} N_k. \quad (4)$$

This statement is a theorem proved in the author's paper [13].

4 The Equation of State

In the simplest version of thermodynamics, one considers the conjugate extensive-intensive pairs pressure-volume, temperature-entropy, chemical potential-number of particles.

One of the main notions of thermodynamics is the equation of state. In 6-dimensional phase space, where the intensive thermodynamical variables P , T , and $-\mu$ play the role of coordinates and the corresponding extensive variables V , $-S$, and N play the role of momenta, the equation of state is described by a 3-dimensional surface on which several additional identities, corresponding to the so-called thermodynamical potentials, hold. This results in the fact that this 3-dimensional surface is a Lagrangian manifold³, and the thermodynamical potential corresponds to action in mechanics.

If one does not consider the number of particles and the volume separately, and only considers their ratio, i.e., density, then one variable turns out to be redundant, and we can consider 4-dimensional phase space and the 2-dimensional Lagrangian surface.

If the volume V is given, we can use a single thermodynamical potential Ω that can be expressed as follows:

$$d\Omega = (-S) dT + N d(-\mu), \quad \Omega = -PV. \quad (5)$$

Besides, since the internal energy is $\tilde{E} = \text{const} \cdot PV$, it is customary in thermodynamical diagrams to plot the pressure P along the abscissa axis, as this was done in the Hougen-Watson diagram.

On the P_r - Z diagram, the *spinodal* is defined as the curve which is the locus of all points where the tangents to the isotherms are perpendicular to the P_r axis (see Figures 2 and 1).

The mathematical model constructed by the author in the papers [1]–[2], which uses the density “effect” mentioned in Section 3 above, shows that

1) if the spinodal in the gaseous region for any gas with non-polarized molecules is given, then all the gas isotherms can be constructed;

2) if we know the slope of the isotherm for $P_r = 0$ on the P_r - Z diagram (this is equivalent to knowing the value of the second virial coefficient [21]) as well as the critical

³In the data base MathSciNet (American Mathematical Society) the query to MathJax on Publications results for “Anywhere =(Maslov index) OR Anywhere (Maslov class)” in a list of 575 papers published before 2013, where the notions “Lagrangian submanifolds” [14], “Maslov index”, “Maslov class” are developed and extended. See also [15]–[16] and so on.

isochore for any fluid (the supercritical state for any collection of non-polarized molecules), then we can construct all the isotherms of the fluid up to the critical isochore;

3) if we know the liquid binodal for any gas consisting of non-polarized molecules, and also know the pressure P_r and the density ρ on the line $Z = 1$ (the Zeno line), then we can construct the liquid isotherms passing through the point $P_r = 0$, $Z = 0$ for so called “hard” liquid consisting of non-polarized molecules on the interval $0 \leq Z \leq 1$.

For an arbitrary liquid, by the *Temperley temperature* we mean the minimal temperature of the set of liquid isotherms passing through the point $P_r = 0$, $Z = 0$. This temperature (corresponding to $K = \infty$), calculated by the physicist Temperley for the Van-der-Waals gas, equals $3^3/2^5 T_c$ [20].

As proved by the author in [1]–[2], the corresponding distribution can be expressed in terms of the polylogarithm⁴

$$\text{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s} = z + \frac{z^2}{2^s} + \frac{z^3}{3^s} + \dots \quad (6)$$

This distribution yields the following expressions for the thermodynamical potential Ω , the density (concentration) ρ and the reduced pressure P_r :

$$\Omega = -\Lambda(\gamma, K) T_r^{2+\gamma} \left\{ \text{Li}_{2+\gamma}(a) - \frac{1}{(K+1)^{1+\gamma}} \text{Li}_{2+\gamma}(a^{K+1}) \right\}, \quad \gamma = \gamma(T_r). \quad (7)$$

$$\rho(T_r, \gamma, K) = \frac{\Lambda(\gamma, K)}{\zeta(2+\gamma_c)} T_r^{1+\gamma} \left(\text{Li}_{1+\gamma}(a) - \frac{1}{(K+1)^\gamma} \text{Li}_{1+\gamma}(a^{K+1}) \right), \quad (8)$$

$$P_r(T_r, \gamma, K) = \frac{\Lambda(\gamma, K)}{\zeta(2+\gamma_c)} T_r^{2+\gamma} \left(\text{Li}_{2+\gamma}(a) - \frac{1}{(K+1)^{1+\gamma}} \text{Li}_{2+\gamma}(a^{K+1}) \right), \quad (9)$$

where K is the maximal occupation number, $a = e^{\mu/T_r}$ (a is called the activity), $\gamma = D/2 - 1$, D is the collective number of degrees of freedom, $\Lambda(\gamma, K)$ is a normalizing constant.

For each type of gas, the critical parameters T_c , P_c , ρ_c are determined experimentally. Substituting these values into (8)–(9), we can determine the parameters $\gamma = \gamma_c$ and $K = K_c$. The parameter γ_c is the number of collective degrees of freedom $D_c = 2\gamma_c + 2$ of the critical isotherm. By K_c we denote the admissible size of clusters (maximal occupation number)⁵ at the energy level of the critical point.

Using (7) and the fact that

$$\tilde{E} = -\mu(\partial\Omega)/(\partial\mu) - T(\partial\Omega)/(\partial T) + \Omega,$$

where $\Omega = -PV$, we derive $\tilde{E} = (D/2)PV$.

5 The notion of generalized ideal gas

We pass to the definition of the new classical ideal gas.

Note that in the description of the behavior of multi-particle systems it is possible that the initial conditions involve interactions. Consider the following analogy. There is

⁴Note that for $z = 1$ and $\text{Re}(s) > 1$, $\text{Li}_s(1) = \zeta(1)$ is Riemann’s zeta function.

⁵The separation of the clusters of molecules near the flat surface of a liquid for $T = T_c$ can be observed experimentally.

a battery of canons. The gunners position the canons and interact with each other in the process. At some moment the command “Fire!” is heard (it means “all interactions (all conversations) must stop”). At the initial situation interactions took place, but once the canon balls are fired, they fly without interacting with each other. However elastic collisions between the balls are possible.

This model can be regarded as a model of ideal gas in the traditional understanding only if the initial data is given at time “minus infinity”.

A generalized classical ideal gas is a classical multi-particle system, where interactions cannot occur only inside certain open regions of the diagram, although at the initial moment of time or on the boundary of a region interactions between particles are allowed. In the region itself, only elastic collisions between particles are possible.

Why is it important to specify the open regions where no interaction (more precisely, no attraction) occurs? Obviously, if one must rapidly mix ball bearings of different sizes, it is easier to do this if the balls are not magnetized. The solution of this problem has several practical applications. Thus the speed with which two gases mix influences the driving force of jet engines: the higher the speed, the higher the driving force. If there is no attraction between molecules of a mixture of gases moving in a pipeline, the probability of the appearance of liquid bottlenecks is considerably less.

Thus the parameters of the Van-der-Waals gas coincide closely enough with the parameters of the new (generalized) gas in the following open regions of the parameters: 1) the region from the critical isotherm ($T_r = 1$) to the critical isochore ($\rho = \rho_c$); 2) the region outside the gas spinodal; 3) the region of hard liquid (below the Temperley temperature, see Section 7).

6 Van-der-Waals forces

The Van-der-Waals model gave rise to the notion of “Van-der-Waals force”. These forces are long-range forces. We will consider the Van-der-Waals forces for the example of the so-called Lennard–Jones potential, which is given by

$$V(r) = \tilde{a} \left(\frac{a^{12}}{r^{12}} - \frac{a^6}{r^6} \right),$$

where $\tilde{a}$ is the depth of the potential well, a is the effective radius, and the attraction is also equal to $1/r^6$. For this potential, long-range Van-der-Waals forces exist.

The quantum dipole-dipole interaction [7], [17] gives an attraction force of the order of $1/r^6$. In the Lennard–Jones potential the attraction is also of the form $1/r^6$. Thus the purely quantum force of attraction smoothly passes into the classical attraction. This was rigorously proved by the author in [18]. But such forces do not exist for all gases. Experiments tell us that particles must be near each other in order to attract each other. This means that attraction is either a quantum effect of dipole-dipole interaction [7], or a Casimir effect, which is also purely quantum. This interaction disappears as $\hbar \rightarrow 0$ (in the classical limit), see [18].

The long-range Van-der-Waals force is true for the Lennard–Jones potential, but certainly it is not true for all potentials. Let us present an analogy. In quantum mechanics, the quasiclassical solution coincides with the exact solution for certain potential, but such potentials are rare, albeit very important.

Thus the theory constructed by the author in [1]–[2] and other papers is not based on “Van-der-Waals forces”, but on combining boundary data with the notion of ideal gas as a gas without interaction of particles with a given number of degrees of freedom.

In other words, in the author’s approach, which is valid for non-polarized molecules, interactions can occur only on the boundary or for densities above the critical level. The critical density is the one above which the molecules are so close to each other (“liquid fluids”) that the Van-der-Waals attracting forces come into play (compare Figures 5 and 10).

7 Comparison of the Generalized Ideal Gas with the Van-der-Waals model

The main practical results that follow from the author’s theory are related to metastable states. In that sense, the main notion of the thermodynamics described here is not the binodal: the curve on the diagram that indicates the stable phase transition “vapor-to-liquid” (the gas binodal) and the phase transition “liquid-to-vapor” (the liquid binodal). The main notion is the spinodal – the curve on the diagram that delimits the regions of stable and metastable states (a more precise definition of the spinodal was given above). It is practically impossible to reach values of the thermodynamical parameters on the spinodal experimentally, because that requires “infinite time”.

From relations (8)–(9) as $K \rightarrow \infty$ we obtain the following equations:

$$P_r = \frac{\Lambda(\gamma) T_r^{2+\gamma} \text{Li}_{2+\gamma}(a)}{\zeta(2+\gamma_c)}, \quad a = e^{\mu/T_r}, \quad \Lambda(\gamma) = \Lambda(\gamma, \infty) = \lim_{K \rightarrow \infty} \Lambda(\gamma, K), \quad (10)$$

$$\rho = \frac{\Lambda(\gamma) T_r^{1+\gamma} \text{Li}_{1+\gamma}(a)}{\zeta(2+\gamma_c)}. \quad (11)$$

For $a = 1$, we obtain the spinodal defined above. This fact was proved by the author in [19].

The constants Λ and γ can be found for $a = 1$ ($\mu = 0$) from the coincidence of the values of P_r and ρ from (10) and (11) with the values of P_{sp} and ρ_{sp} on the Van-der-Waals spinodal.

In Figures 3 and 4, the isotherms of the Van-der-Waals model are compared with those of the author’s model on Hougen–Watson diagrams.

In Figure 4, it is shown how the isotherms of the ideal gas (in the new sense) obtained from (10)–(11), differ from the isotherms of the Van-der-Waals model. The figure shows that the isotherms practically coincide except at points near critical values. In regions with less than critical parameter values, the coincidence is up to 0.006%, while at points near critical values the error is of the order of 0.01%. Note the important point $0.9T_c$. At that point, the maximal discrepancy between the isotherms is observed. It is of the order of ~ 0.01 .

For isotherms corresponding to temperatures less than the Temperley temperature (“cluster-free liquids”), we obtain a very precise coincidence between the isotherms constructed according to the distribution (10)–(11) and the isotherms coming from the Van-der-Waals model. The temperature $T_{\text{Temperley}}$ specifies a boundary above which the number K (maximal occupation number) becomes infinite.




Figure 3: On this diagram the pressure is $P = P_r$. Coincidence of the Van-der-Waals isotherms (solid lines) with the isotherms from the author's model (dashed lines). The following isotherms are shown $T_r = 0.1$, $T_r = 0.2$, $T_r = 0.3$, $T_r = 0.4$, $T_r = 0.5$, $T_r = 0.6$, $T_r = 0.7$, $T_r = 0.8$, $T_r = 0.9$, $T_r = 1$. The value of $\gamma = \gamma(T_r)$ in equations (10)–(11) is determined from the relations on the Van-der-Waals spinodal $Z = Z_{sp}$, $P = P_{sp}$ for $a = 1$.




Figure 4: Discrepancy between the isotherms of the Van-der-Waals model and the isotherms obtained from the new distribution; $a = e^{\mu/T_r}$ is the activity, T_r is the reduced temperature. The maximal discrepancy is ~ 0.01 for $T_r = 0.9$ (the effect of $K \neq \infty$ is substantial).

8 Supercritical Energy

We introduce an additional parameter, the integer K , specifying the maximal occupation number for clusters when particles do not interact with each other except at the boundaries of the regions. First of all, this is the open region of ordinary ideal gas, where the density vanishes $\rho = 0$. The boundary conditions are well determined by the second virial coefficient, which can be found from the Van-der-Waals equation.. This coefficient, as we indicated above, gives the slope of the isotherm on the Hougen–Watson diagram [21].

If we consider the situation for temperatures above the Temperley temperatures, then we have to specify one more constant, i.e., the initial conditions on the spinodal do not suffice. This second condition can also be found from the critical isochore of the Van-der-Waals gas. We are dealing with ideal gas, i.e., a gas without interaction (attraction) of particles for $\rho = 0$. However, in those regions where the density is high, i.e., $\rho \geq \rho_c$, attraction between particles can occur. It is on the isochore $\rho = \rho_c$ that we must take the second initial condition.

So what we are doing is imposing boundary conditions on the derivative on one of the

boundaries and imposing ordinary boundary conditions on the other.




Figure 5: On this plot $P = P_r$. The Van-der-Waals isotherms (solid lines) and the isotherms from the author's theory corresponding to formulas (8)–(9) (hashed lines) for $T_r = 1$, $T_r = 1.2$, $T_r = 1.5$, $T_r = 1.8$.

Simply speaking, the boundary means that for “time” $t < t_0$ the balls interacted, then for $t = t_0$ the interaction vanished (the role of time here is played by the chemical potential μ). On the other boundary, one can find two constants: the number of degrees of freedom and the maximal occupation number K . After that we can compare the isotherms of the ideal parastatistical (noninteracting) gas⁶ with the corresponding Van-der-Waals gas isotherms. See Figure 5.

9 Liquids

Now let us consider liquids.

In liquids collisions of molecules occur so often that they limit the number of collective degrees of freedom of the molecules to two and hamper their arbitrary motion. The system of prohibitions for the collection of molecules can be considered in a more detailed and complicated way in the framework of their quantum-mechanical description (see [22]).

We will call *congestion* any state of a system of molecules for which the collective number of degrees of freedom is less than two. The liquids that we consider are systems of molecules in the state of congestion.

Let us consider some analogies. In a normal state, an individual locust has 3 degrees of freedom (in the sense that it can move in three independent directions). The uniting of locusts into a dense flock is similar to the phase transition of gas into liquid [23]. Another analogy has to do with human beings. A person in a crowd is subjected to a number of exclusions related to the people near him and to the psychology of crowds.

The above analogies correspond to the new notion of liquid consisting of noninteracting particles (balls) with less than two degrees of freedom – a generalization of the notion of hard liquid. A very similar model of liquid without attraction as a dense incompressible ideal gas was considered earlier by some physicists [24]–[25].

The value $\gamma = 0$ (or, what is the same, the value 2 for the number D of degrees of freedom) is in a certain sense critical. The author proved in [26], that if $\gamma < 0$, the spectrum near zero becomes very sparse (compare (2)). This means that a *gap* is formed between the zero energy level and the energy level nearest to it.

⁶Such a gas is analogous to the noninteracting canon balls mentioned above, which can collide but still remain similar to particles of an ideal gas.

This situation shows that the density (concentration) of particles on the zero energy level does not depend on the chemical potential μ . This density for the Van-der-Waals model can be calculated as a function of temperature:

$$\rho_0(T_r) = \frac{3}{2} + \sqrt{\frac{9}{4} - \frac{8T_r}{3}}. \quad (12)$$




Figure 6: On this plot, $P = P_r$. The isotherm here is $T = T_{\text{Temperly}} = 3^3/2^5 T_c$. The solid line is the Van-der-Waals isotherm, the hashed line is the isotherm of the generalized ideal gas with two degrees of freedom ($\gamma = 0$), the dotted line is the straight line joining the point $P_r = 0, Z = 0$ to the point $T = T_{\text{Temperly}}$ on the Zeno line.




Figure 7: On this plot $P = P_r$. There are three isotherms with $T_r = 0.4$, $T_r = 0.6$, $T_r = 0.8$. The solid lines are the Van-der-Waals isotherms, the hashed lines are the isotherms from the author's model, the dotted lines are straight lines passing through the point $P_r = 0, Z = 0$ and the corresponding point on the Zeno line (i.e., on the line $Z = 1$).

The “theoretical” curves shown in Figures 6 and 7 were constructed by using the relations derived in the author's model

$$\rho \equiv -\frac{\tilde{\Lambda}(\gamma)}{\zeta(2 + \gamma_c)} T_r^{1-\gamma} \text{Li}_{1+\gamma}(-a) + \rho_0(T_r), \quad (13)$$

$$P_r = -\frac{\Lambda(\gamma)}{\zeta(2 + \gamma_c)} T_r^{2-\gamma} \text{Li}_{2+\gamma}(-a), \quad (14)$$

where $\gamma \leq 0$ and the polylogarithms are defined by (6), as well as by the following integral formula

$$-\text{Li}_s(-z) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1}}{e^t/z + 1} dt. \quad (15)$$

In order to find all unknown constants needed to determine the “theoretical” liquid isotherms, we must satisfy two boundary conditions by choosing the liquid binodal curve and the Zeno line $Z = 1$. For the liquid binodal, we have

$$\begin{aligned} P_r|_{\text{bin}} &= \frac{\Lambda(\gamma)}{\zeta(2 + \gamma_c)} T_r^{2+\gamma} (-\text{Li}_{2+\gamma}(-e^{\mu_{\text{bin}}/T_r})), \\ \rho_{\text{bin}} &= \rho_0(T_r) + \frac{\tilde{\Lambda}(\gamma)}{\zeta(2 + \gamma_c)} T_r^{1+\gamma} (-\text{Li}_{1+\gamma}(-e^{\mu_{\text{bin}}/T_r})). \end{aligned} \quad (16)$$

Similar relations for $P_r|_{\text{Zeno}}$ and ρ_{Zeno} have the form

$$\begin{aligned} P_r|_{\text{Zeno}} &= \frac{\Lambda(\gamma)}{\zeta(2 + \gamma_c)} T_r^{2+\gamma} (-\text{Li}_{2+\gamma}(-e^{\mu_{\text{Zeno}}/T_r})), \\ \rho_{\text{Zeno}} &= \rho_0(T_r) + \frac{\tilde{\Lambda}(\gamma)}{\zeta(2 + \gamma_c)} T_r^{1+\gamma} (-\text{Li}_{1+\gamma}(-e^{\mu_{\text{Zeno}}/T_r})). \end{aligned} \quad (17)$$

Relations (16)–(17) produce the plots shown in Figures 6 and 7.

The value ρ_0 remains the same of all values of the chemical potential μ .

All the isotherms of the liquid phase of the Van-der-Waals model for temperatures below the Temperley temperature ($0.84T_c$) pass through the point $Z = 0$, $P_r = 0$ on the Hougen–Watson diagram, which is a node for all the isotherms, and so, for $T_r > 0.84T_c$ (more precisely, for $T_r \geq (3^3/2^5)T_c$) the isotherms are those of a “hard” liquid.

10 Concluding Remarks: Quasi-statical Processes and the Two-Energy Dispersive Model

Caratheodory proposed an approach in which thermodynamics is described as a “limiting” theory. That approach is very important because it indicates the directions in which this “limiting” model can be extended.

Caratheodory introduced the notion of equilibrium quasi-statical process in thermodynamics; in such a process, equilibrium is achieved by damping forces such as friction (viscosity), after which the damping forces are considered negligible. His idea is that the damping forces (viscosity) are needed to achieve equilibrium, and after that must be disregarded, so that an invertible process (a quasi-statical process) begins.

As shown by the author in [27], the definition of quasi-statical process given by Caratheodory requires additional constraints or additional properties that consist in the following. At the beginning of a quasi-statical process, besides the damping effects, an important role is played by the gravitational attraction of the Earth. This attraction is small, but nevertheless it plays its role in bringing the process to a state that may be described in the framework of phenomenological thermodynamics.

If we try to carry out this procedure with mathematical accuracy, we must first argue in the framework of hydrodynamics, where viscosity occurs, and, after a long period of time, make viscosity tend to zero. But it turns out that this can’t always be done. It cannot be done on the spinodal, in particular, near the critical point. In a small region near the critical point (in which the temperature differs from the critical one by ± 0.005 degrees or less) a phenomenon akin to turbulence sets in. This phenomenon is called *opalescence* because besides turbulence and vortex effects, a change in the color of light takes place.

Let us return to the role of Earth's gravitation. The so-called *dew point* is the state of the gas in which little droplets (tiny balls) appear (they will form what in everyday language is called dew). These little balls are not stable until their radius reaches a certain critical value. In that situation the pressure inside the droplet will be greater than the pressure outside it. If additional relations due to surface tension are not introduced, then we come to a contradiction with one of the most important laws of equilibrium. In order to overcome this contradiction, sometimes additional equations are given, thus making the mathematical model more complicated.

But once we take the Earth's gravitation into account, the following picture arises. The amalgamation of droplets leads to an energetically more favorable state. Hence all the droplets, falling to the bottom of the test tube, will merge, and an almost flat boundary of the water appears (provided the test tube is not too narrow). We then obtain the correct "gas-liquid" equilibrium relations that were formulated in the past by the great physicists. Therefore we must take into account the long period of time needed for the droplets to reach the bottom. And so the Earth's gravitation must play its role in this process as well.

The role of the Earth's gravitation is also important in the following situation. When heaters are turned on in a house, after a while equilibrium will be reached and a uniform temperature will be established in the house, so that the temperature in different rooms will be approximately the same. However, this happens because of *convection*: warm air from the heater will rise because of the laws of gravitation (the Earth's attraction). Whirling flows of hot air circulate and finally, because of viscosity, equilibrium sets in.

The fact that we consider a process which is limiting with respect to viscosity allows the introduction of corrections that take into account viscosity near the critical point. Similarly, by taking into account Earth's gravitation, we can put the clusters into a certain order when $K < K_c$ (where K_c is the maximal occupation number at the energy level of the critical point)⁷. This involves the focal points as well as turbulence; see [28]–[30].

For $\rho > \rho_c$, the molecules are so close to each other that attraction arises. In this case, the particle interaction is described by the double well potential [26] and the family $\{\varepsilon_k\}$ of discrete energy levels in Eq. (2) splits into two series – one with a minimal (zero) energy level and the other with positive values of minimal energy. Particles with energies from the second series move with relatively high velocities even when their energy is near the minimum energy for that series. The mean number of particles in each series can be determined. The mean energy corresponds to the mean value of all the energy levels and includes both series (see also [26]).

As the result, a picture similar to the Tissa–Landau two-liquid model arises; in it, the superfluid component flows through the normal liquid without any resistance. For this reason, the second series can be called superfluid. In our case, this is not a quantum effect, it is the consequence of the Caratheodory axioms, leading to an invertible process, which require that we put viscosity equal to zero. It is similar to the "superfluid" motion of gas without resistance.

As the author has proved, for $\rho > \rho_c$, the supercritical fluid splits into two series: a cluster sponge, consisting of clusters of liquid type and gas monomers with viscosity equal to zero that interact with each other, see [26], [31]. Obviously, in such a two-fluid system, the pressure in both components is the same. Hence the distribution of pressure does not change, while the spectrum will accumulate near zero as density increases because of the

⁷ The separation of clusters of molecules with a certain occupation number near the flat surface of a liquid for $T = T_c$ can be observed experimentally.




Figure 8: The double well potential $E = u(x)$. The normal series and the superfluid one for energies $\{\varepsilon_k\}$ in equation (2).

gravitational attraction of Earth. By comparing to Van-der-Waals gas, we find the value of the cut-off parameter A (from momenta) of the form (18) for each point of the isotherm.




Figure 9: Values of the parameter $A(P_r, T_r)$ in the Van-der-Waals model for $T_r = 1$, $T_r = 1.2$, $T_r = 1.5$, $T_r = 1.8$ for $\rho > \rho_c$. On the plot $P = P_r$.

Therefore the integrals appearing [5] which give the distribution of momenta (for energy) must be cut off to the value A :

$$\rho(T_r, \gamma, K) = \frac{\Lambda(\gamma, K)}{\zeta(2 + \gamma_c)} \int_0^{A(P_r, T_r)} \left(\frac{1}{e^{(x-\mu)/T_r} - 1} - \frac{K+1}{e^{(K+1)(x-\mu)/T_r} - 1} \right) x^\gamma dx, \quad (18)$$

$$P_r(T_r, \gamma, K) = \frac{\Lambda(\gamma, K)}{\zeta(2 + \gamma_c)} \int_0^\infty \left(\frac{1}{e^{(x-\mu)/T_r} - 1} - \frac{K+1}{e^{(K+1)(x-\mu)/T_r} - 1} \right) x^{1+\gamma} dx, \quad (19)$$

where $-1 < \gamma < \infty$.

The value of the cut-off parameter $A(P_r, T_r)$ is the smaller when the density of the normal series is the greater – because of the law of the Earth’s attraction. The parameter A may be chosen so that the isotherms (18)-(19) will coincide with the isotherms of the Van-der-Waals isotherms, see Figure 9. Naturally, the pressure does not depend on the series.

Figure 10 shows the coincidence of the isotherms in the Van-der-Waals model with the isotherms obtained by the author for the case where the value of $A(P_r, T_r)$ is chosen due to the Van-der-Waals model.




Figure 10: Coincidence of the Van-der-Waals isotherms (solid lines) with the isotherms obtained for $A = A(P_r, T_r)$ (dotted lines) for the temperatures $T_r = 1$, $T_r = 1.2$, $T_r = 1.5$, $T_r = 1.8$. On this diagram $P = P_r$. The interacting region $\rho \geq \rho_c$ corresponds to the two-energy dispersive model.

Thus, by taking into account the Earth's gravitational attraction, we split the fluid obtained when $\rho > \rho_c$ into two dispersive regions. Sometimes experimenters actually see a surface film, as in the gas-liquid picture. To justify this model, the law of conservation of the number of degrees of freedom and the Tissa–Landau model are used.

Remark 2. At $T_r \sim 0.5$ liquids turn into solids. This fact is not taken into account in the Van-der-Waals model. Therefore the comparison of the isotherms of the Van-der-Waals model with the isotherms derived by the author can only be made near the Temperley temperature [32], see Figure 11.




Figure 11: The Van-der-Waals isotherm $T = 0.4$ (solid line) and the isotherm of the author's model (dashed line), where the activity a in equations (13)–(14) changes sign at the origin $P_r = 0$, $Z = 0$. On this diagram $P = P_r$.

The author is deeply grateful to professor V.S. Vorobiev for fruitful discussions as well as for checking all the diagrams and verifying the author's model on the basis of the latest experimental results for nitrogen. The author is also grateful to E.P. Loubenets for useful discussions and D.S. Minenkov for help in constructing the diagrams.

References

- [1] Maslov V. P. Undistinguishing Statistics of Objectively Distinguishable Objects: Thermodynamics and Superfluidity of Classical Gas. Math. Notes 2013; **94** (5):722–813.

- [2] Maslov V. P. New Construction of Classical Thermodynamics and UD-Statistics. Russian J. Math. Phys. 2014; **21** (2):256-284.
- [3] Maslov V. P. Thermodynamics and Economics: Overview. Reference Module in Earth Systems and Environmental Sciences. Elsevier; 2014.
- [4] Dobrokhotoy S. Yu., Karasev M. V., Kolokoltsov V. N., et al. Award of the State Prize of the Russian Federation in Science and Technology (2013) to Victor Maslov. Probability Theory and Applications 2014; **59** (2):2009–2013.
- [5] Landau L. D. and Lifshits E. M. Statistical Physics. Moscow: Nauka; 1964 [in Russian].
- [6] Kadomtsev B. B. Dynamics and Information. Moscow, Uspekhi Fizicheskikh Nauk; 1999 [in Russian].
- [7] Landau L. D. and Lifshits E. M. Quantum Mechanics: Non-Relativistic Theory, 2nd ed. Moscow:Nauka; 1964 [in Russian]; (translation of the 1st ed., Pergamon Press, London–Paris and Addison-Wesley Publishing Co., Inc., Reading, Mass.; 1958).
- [8] Maslov V. P. and Shvedov O. Yu. The Complex Germ Method in Many-Particle Problems and in Quantum Field Theory. Moscow: Editorial URSS; 2000 [in Russian].
- [9] Dai W.-S., Xie M. Gentile statistics with a large maximum occupation number. Annals of Physics 2004 **309**:295–305.
- [10] Maslov V. P. The relationship between the Van-der-Waals model and the undistinguishing statistics of objectively distinguishable objects. The new parastatistics. Russian J. Math. Phys. 2014; **21** (1):99–111.
- [11] Vershik A. M. Statistical mechanics of combinatorial partitions, and their limit shapes,” Funktsional. Anal. i Prilozhen. 1996; **30** (2): 19–39 [Functional Anal. Appl. 1996; **30** (2):90–105].
- [12] Maslov V. P. Nazaikinskii V. E. On the rate of convergence to the Bose-Einstein distribution. Math. Notes 2016; **99** (1):95–109.
- [13] Maslov V. P. Case of less than two degrees of freedom, negative pressure, and the Fermi–Dirac distribution for a hard liquid,” Math. Notes 2015; **98** (1):138–157.
- [14] Maslov V. P. Perturbation Theory and Asymptotical Methods. Moscow: Izd. Moskov. Univ.;1965; Paris: Dunod;1972 [in Russian and French].
- [15] Bayfield J. E. Quantum Evolution. New York–Toronto: John Wiley and Sons; 1999.
- [16] Römer H. Theoretical Optics. Wienheim: Wiley-VCH; 2005.
- [17] Maslov V. P., Fedoriuk M. V. Semi-Classical Approximation in Quantum Mechnics. Dordrecht-Boston-London: D. Reidel Publishing Company; 1981.
- [18] Maslov V. P. On the Van-der-Waals forces. Math. Notes 2016; **99** (2):284–289.
- [19] Maslov V. P. Gas–amorphous solid and liquid–amorphous solid phase transitions. Introduction of negative mass and pressure from the mathematical viewpoint. Math. Notes 2015; **97** (3):423–430.

- [20] Temperly H. N. // Proc. Phys. Soc. (L.) 1947; **59**:199–205.
- [21] Burshtein A. I. Molecular Physics. Novosibirsk: Nauka; 1986 [in Russian].
- [22] Percival I. C. Semiclassical theory of bound states. Advances in Chemical Physics, **36**, edited by Ilya Prigogine and Stuart A. Rice. New-York-London-Sydney-Toronto: John Wiley and Sons; 1977: 1–62.
- [23] Maslov V. P. Theory of chaos and its application to the crisis of debts and the origin of the inflation. Russian J. Math. Phys. 2009; **16** (1):103–120.
- [24] Anselm A. I. Foundations of Statistical Physics and Thermodynamics. Moscow: Nauka; 1973 [in Russian].
- [25] Eyring H. J., Lin S.H., Lin S. M. Basic chemical kinetics. New York: John Wiley and Sons Inc; 1980.
- [26] Maslov V. P. Two-fluid picture of supercritical phenomena. Teoret. Mat. Fiz. 2014; **180** (3):1095–1128. [Theoret. and Math. Phys. 2014; **180** (3):1095–1128].
- [27] Maslov V. P. “Violation of Carathéodory axioms at the critical point of a gas. Frenkel point as a critical point of the transition “liquid-amorphous solid” in the region of negative pressure,” Math. Notes **96** (6), 977–982 (2014).
- [28] Danilov V. G., Maslov V. P. Volosov K. A. Mathematical Modelling of Heat and Mass Transfer Processes. Dordrecht / Boston / London: Kluwer Academic Publishers; 1995.
- [29] Maslov V. P. Kolmogorov’s law and the Kolmogorov scales in anisotropic turbulence: the occurrence of turbulence as a result of three-dimensional interaction. Teoret. Mat. Fiz. 1993; **94** (13):368–374.
- [30] Maslov V. P., Shafarevich A. I. Asymptotic solutions of Navier-Stokes equations and topological invariants of vector fields and Liouville foliation. Theoretical and Math. Physics, 2014; **180** (2):967–982.
- [31] Maslov V. P. Calculation of the number of collective degrees of freedom and of the admissible cluster size for isotherms in the Van-der-Waals model in supercritical states. Russian J. Math. Phys. 2014; **21** (4):494–503.
- [32] Maslov V. P., Maslova T. V. New thermodynamics and frost cleft in conifers. Math. Notes 2015; **98** (2):343–347.